arXiv:math/0406590v1 [math.OA] 29 Jun 2004

TOPOLOGICAL ENTROPY AND AF SUBALGEBRAS OF GRAPH C^*-ALGEBRAS

JA A JEONG[†] AND GI HYUN PARK[‡]

Abstract. Let $\mathcal{A}_E$ be the canonical AF subalgebra of a graph C^*-algebra $C^*(E)$ associated with a locally finite directed graph E. For Brown-Voiculescu's topological entropy $ht(\Phi_E)$ of the canonical completely positive map Φ_E on $C^*(E)$, $ht(\Phi_E) = ht(\Phi_E|_{\mathcal{A}_E}) = h_l(E) = h_b(E)$ is known to hold for a finite graph E, where $h_l(E)$ is the loop entropy of Gurevic and $h_b(E)$ is the block entropy of Salama. For an irreducible infinite graph E, the inequality $h_l(E) \le ht(\Phi_E|_{\mathcal{A}_E})$ has been known recently. It is shown in this paper that

$$ht(\Phi_E|_{\mathcal{A}_E}) \le \max\{h_b(E), h_b({}^tE)\},$$

where ${}^tE$ is the graph E with the direction of the edges reversed. Some Irreducible infinite graphs $E_p (p > 1)$ with $ht(\Phi_E|_{\mathcal{A}_{E_p}}) = \log p$ are also examined.

1. Introduction

Voiculescu [20] introduced a notion of topological entropy $ht(\alpha)$ for an automorphism α of a nuclear unital C^*-algebra A to measure the growth of α^n as $n \to \infty$ using the fact that a nuclear C^*-algebra has the completely positive approximation property. The definition extends very well to automorphisms of exact C^*-algebras (as done by Brown in [4]) due to the deep result by Kirchberg [12] that exact C^*-algebras are nuclearly embeddable. But without effort one can define $ht(\Phi)$ even for a completely positive (cp) map on an exact C^*-algebra as described in [2]. Since a C^*-subalgebra of an exact C^*-algebra is always exact, if $\Phi : A \to A$ is a cp map on an exact C^*-algebra A and B is a Φ-invariant C^*-subalgebra of A then $ht(\Phi|_B)$ can be defined and the monotonicity $ht(\Phi|_B) \le ht(\Phi)$ holds.

The topological entropy has been computed in several cases, for example, the equality $ht(\alpha * \beta) = \max\{ht(\alpha), ht(\beta)\}$ for the reduced free product automorphism $\alpha * \beta$ was proved in [1], when the free product is with amalgamation over a finite dimensional C^*-algebra. Also Dykema [9] showed that $ht(\alpha) = 0$ for certain classes of automorphisms α of reduced amalgamated free products of C^*-algebras, which turns out to extend Størmer's result [19] that the Connes-Størmer entropy of the free shift automorphism of the II_1-factor $L(F_\infty)$ is zero.

In this paper we are concerned with the topological entropy of the shift type cp maps on C^*-algebras arising from directed graphs. A typical one is the canonical cp map $\Phi_A : \mathcal{O}_A \to \mathcal{O}_A$ of the Cuntz Krieger algebra $\mathcal{O}_A$ given by

$$\Phi_A(x) = \sum_{i=1}^{n} s_i x s_i^*,$$

2000 *Mathematics Subject Classification.* 46L05, 46L55.

[†] Partially supported by KOSEF R14-2003-006-01000-0.

[‡] Partially supported by KOSEF R01-2001-000-00001-0.

where $s_1, \dots, s_n$ are the partial isometries that generate $\mathcal{O}_A$. The reason we call Φ_A shift type is that $\mathcal{O}_A$ contains a Φ_A-invariant commutative C^*-subalgebra $\mathcal{D}_A$ which is isomorphic to $C(X_A)$ in such a way that the restriction $\Phi_A|_{\mathcal{D}_A}$ corresponds to the shift map σ_{X_A} on the (compact) shift space X_A associated with the incidence matrix A. The topological entropy of Φ_A is then computed (see [5], [2], [17]) as $ht(\Phi_A) = \log r(A)$ ($r(A)$ is the spectral radius of A). But $\log r(A) = h_{top}(X_A)$ is a well known fact, so that one can deduce by [8] that $ht(\Phi_A) = ht(\Phi_A|_{\mathcal{D}_A})$. On the other hand, $\mathcal{O}_A$ also contains another important Φ_A-invariant C^*-subalgebra $\mathcal{A}_A$ which is an AF algebra with $\mathcal{D}_A \subset \mathcal{A}_A$. Thus by monotonicity of entropy, we have $ht(\Phi_A) = ht(\Phi_A|_{\mathcal{A}_A}) = ht(\Phi_A|_{\mathcal{D}_A})$.

The Cuntz-Krieger algebras $\mathcal{O}_A$ are now well understood as graph C^*-algebras $C^*(E) = C^*(s_e, p_v)$ associated with finite directed graphs E and the cp map Φ_A of $\mathcal{O}_A$ is interpreted as the map $\Phi_E : C^*(E) \to C^*(E)$ given by $\Phi_E(x) = \sum_{e \in E^1} s_e x s_e^*$. Hence if E is a finite graph (possibly with sinks) which contains an infinite path, it follows that $ht(\Phi_E) = ht(\Phi_E|_{\mathcal{A}_E}) = ht(\Phi_E|_{\mathcal{D}_E}) = \log r(A_E)$, where $\mathcal{A}_E$ is the AF subalgebra of $C^*(E)$ corresponding to $\mathcal{A}_A$ in $\mathcal{O}_A$ and A_E is the edge matrix of E (see [11]).

If E is infinite but locally finite then the map Φ_E is known to be a contractive cp map, and furthermore if E is irreducible and $\mathcal{A}_E$ is the canonical AF subalgebra of $C^*(E)$, the inequality $h_l(E) \le ht(\Phi_E|_{\mathcal{A}_E})$ is known to hold [11]. The purpose of the present paper is then to give an upper bound for $ht(\Phi_E|_{\mathcal{A}_E})$ and we actually prove the following (see Theorem 3.9)

$$ht(\Phi_E|_{\mathcal{A}_E}) \le \max\{h_b(E), h_b({}^tE)\}.$$

In particular, for an irreducible infinite graph E_p constructed in [18] so that $h_l(E_p) = h_b(E_p) = p > 1$, we have $ht(\Phi_{E_p}|_{\mathcal{A}_{E_p}}) = \log p$.

We believe that the result would be helpful to compute the entropy $ht(\Phi_E)$ of Φ_E on the whole graph C^*-algebra $C^*(E)$.

2. Preliminaries

2.1. **Graph C^*-algebras.** A (directed) graph is a quadruple $E = (E^0, E^1, r, s)$ of the vertex set E^0, the edge set E^1, and the range, source maps $r, s : E^1 \to E^0$. A family $\{p_v, s_e \mid v \in E^0, e \in E^1\}$ of mutually orthogonal projections p_v and partial isometries s_e is called a *Cuntz-Krieger E-family* if the following relations hold;

$$\begin{aligned} & s_e^* s_e = p_v, \quad s_e s_e^* \le p_{s(e)}, \\ & p_v = \sum_{s(e)=v} s_e s_e^*, \quad \text{if } 0 < |s^{-1}(v)| < \infty. \end{aligned}$$

The *graph C^*-algebra $C^*(E)$* is then defined to be a C^*-algebra generated by a universal Cuntz-Krieger E-family (see [15], [3]). We call E *locally finite* if each vertex receives and emits only finitely many edges. Throughout this paper we consider only locally finite graphs, and adopt the notations in [15]. If a finite path $\alpha \in E^*$ of length $|\alpha| > 0$ is a return path, that is, $s(\alpha) = r(\alpha)$, then α is called a *loop* at $v = s(\alpha)$. A graph E is said to be *irreducible* if for any two vertices v, w there is a finite path $\alpha \in E^*$ with $s(\alpha) = v$ and $r(\alpha) = w$. It is known that if E is irreducible and every loop has an exit then $C^*(E)$ is simple.

2.2. **Topological entropy of cp maps.** Let A be a C^*-algebra, $\pi : A \to B(H)$ a faithful $*$-representation, and $Pf(A)$ be the set of all finite subsets of A. For

$\omega \in Pf(A)$ and $\delta > 0$, put

$$\begin{aligned} CPA(\pi, A) &= \{(\phi, \psi, B) \mid \phi : A \to B, \psi : B \to B(H) \text{ are contractive cp maps} \\ &\qquad \text{and } \dim B < \infty\}, \\ rcp(\pi, \omega, \delta) &= \inf\{rank(B) \mid (\phi, \psi, B) \in CPA(\pi, A),\ \|\psi \circ \phi(x) - \pi(x)\| < \delta, \\ &\qquad \text{for all } x \in \omega\}, \end{aligned}$$

where $rank(B)$ denotes the dimension of a maximal abelian subalgebra of B.

Since the cp δ-rank $rcp(\pi, \omega, \delta)$ is independent of the choice of π (see [4], [2]) and graph C^*-algebras $C^*(E)$ are nuclear we may write $rcp(\omega, \delta)$ for $rcp(\pi, \omega, \delta)$ assuming that $C^*(E) \subset B(H)$ for a Hilbert space H.

Definition 2.1. ([4], [2]) Let $A \subset B(H)$ be a C^*-algebra and $\Phi : A \to A$ be a cp map. Put

$$\begin{aligned} ht(\Phi, \omega, \delta) &= \limsup_{n\to\infty} \frac{1}{n} \log \left(rcp(\cup_{i=0}^{n-1} \Phi^i(\omega), \delta) \right) \\ ht(\Phi, \omega) &= \sup_{\delta > 0} ht(\Phi, \omega, \delta). \end{aligned}$$

Then $ht(\Phi) := \sup_{\omega \in Pf(A)} ht(\Phi, \omega)$ is called the *topological entropy* of Φ.

Remark 2.2. We refer the reader to [2] and [4] for the following useful properties and their proofs. Let A be an exact C^*-algebra and $\Phi : A \to A$ be a cp map.

(a) If $\theta : A \to B$ is a C^*-isomorphism then $ht(\Phi) = ht(\theta \Phi \theta^{-1})$.
(b) Let $\tilde{A}$ be the unital C^*-algebra obtained by adjoining a unit and $\tilde{\Phi} : \tilde{A} \to \tilde{A}$ be the extension of Φ. Then $ht(\tilde{\Phi}) = ht(\Phi)$.
(c) If $A_0 \subset A$ is a Φ-invariant C^*-subalgebra of A, $ht(\Phi|_{A_0}) \leq ht(\Phi)$.

We will use the following Arveson's extension theorem several times.

Arveson Extension Theorem (see [4]) Let A be a unital C^*-algebra, $S \subset A$ a unital subspace with $S = S^*$, and $\phi : S \to B$ be a contractive cp map where $B = B(H)$ or $dim(B) < \infty$. Then ϕ extends to a cp map $\bar{\phi} : A \to B$. If S is a C^*-subalgebra of A then we obtain a unital cp extension of ϕ even when S does not contain the unit of A.

If E is a locally finite graph, the map $\Phi_E : C^*(E) \to C^*(E)$, defined by

$$\Phi_E(x) = \sum_{e \in E^1} s_e x s_e^*,$$

is well defined, contractive, and completely positive [11]. For a finite graph E, the topological entropy $ht(\Phi_E)$ has been obtained as follows (see [2], [5], [17], or [11]).

Theorem 2.3. *Let E be a finite graph possibly with sinks and A_E be the edge matrix of E. If E contains an infinite path then*

$$ht(\Phi_E) = \log r(A_E),$$

where $r(A_E)$ is the spectral radius of A_E.

By $h_{top}(X)$ we denote the topological entropy of a compact space (X, T) together with a continuous map $T : X \to X$ (for definition, see [16, Definition 4.1.1] or [13, p.23]). Let E be a locally finite infinite graph and X_E the locally compact shift space of (one-sided) infinite paths with the one point compactification $\bar{X}_E$. Then the first identity in the following theorem is shown for the doubly infinite path space of E by Gurevic [10]. See Definition 3.1 for $h_b(E)$.

Theorem 2.4 ([11, Theorem 4.4]). *Let E be a locally finite irreducible infinite graph. Then*

$$h_{top}(\bar{X}_E) = \sup_{E'} h_b(E') \ \le \ ht(\Phi_E),$$

where the supremum is taken over all the finite (irreducible) subgraphs of E.

3. Main results

Throughout this section E will denote a locally finite infinite graph unless stated otherwise. For a path $\alpha \in E^*$, let α^0 be the set of vertices lying on $\alpha = \alpha_1 \cdots \alpha_n$, that is, $\alpha^0 = \{s(\alpha_1), r(\alpha_1), \ldots, r(\alpha_n)\}$. For a fixed vertex v we consider the following subsets of finite paths E^n of length n.

(i) $E^n(v) = \{\alpha \in E^n \mid v \in \alpha^0\}$,
(ii) $E^n_s(v) = \{\alpha \in E^n \mid s(\alpha) = v\}$,
(iii) $E^n_s(v^\star) = \{\alpha \in E^n_s(v) \mid r(\alpha_i) \neq v,\ 1 \le i \le n\}$,
(iv) $E^n_l(v) = \{\alpha \in E^n \mid \alpha \text{ is a loop at } v\}$.

Similarly we can think of $E^n_r(v)$ and $E^n_r(v^\star)$.

Definition 3.1. Let E be an irreducible graph and let $v \in E^0$.

(a) $h_l(E) := \limsup_n \frac{1}{n} \log |E^n_l(v)|$ is called the *loop entropy* of E.

(b) $h_b(E) := \limsup_n \frac{1}{n} \log |E^n_s(v)|$ is called the *block entropy* of E.

Note that both entropies $h_l(E)$ and $h_b(E)$ are independent of the choice of a vertex v [18]. If ${}^tE$ denotes the graph E with the direction of all edges reversed, then clearly $h_l(E) = h_l({}^tE)$ while $h_b(E) \neq h_b({}^tE)$ in general as we will see in Example 3.3.

We will use the following notation for the infinite series with coefficients from (i)-(iv) above.

(i)′ $E(v,z) := \sum |E^n(v)| z^n$,
(ii)′ $E_s(v,z) := \sum |E^n_s(v)| z^n$,
(iii)′ $E_s(v^\star,z) := \sum |E^n_s(v^\star)| z^n$,
(iv)′ $E_l(v,z) := \sum |E^n_l(v)| z^n$.

We denote the radius of convergence of the series $E_s(v^\star, z)$ by $R_{E^\star_s}$. Thus

$$R^{-1}_{E^\star_s} = \limsup_{n\to\infty} \left|E^n_s(v^\star)\right|^{1/n}.$$

Similarly $R_{E^\star_r}$ denotes the radius of convergence of $E_r(v^\star, z) := \sum |E^n_r(v^\star)| z^n$.

Proposition 3.2 ([18]). *If E is an irreducible graph, then*

$$h_b(E) = \max\{\log\left(R^{-1}_{E^\star_s}\right),\ h_l(E)\}.$$

Note that if E is irreducible then $h_b({}^tE) = \limsup \frac{1}{n} \log |E^n_r(v)|$ and so from the above proposition we have

$$h_b({}^tE) = \max\{\log\left(R^{-1}_{E^\star_r}\right),\ h_l(E)\}.$$

The following example shows that $h_b(E) \neq h_b({}^tE)$ in general.

Example 3.3. For each pair of positive real numbers $1 < p \leq q$, Salama [18] constructed an irreducible infinite graph $E_{p,q}$ with

$$h_l(E_{p,q}) = \log p \quad \text{and} \quad h_b(E_{p,q}) = \log q.$$

For example, the following graph $E := E_{2,8}$ satisfies $h_l(E) = \log 2$ and $h_b(E) = \log 8$. There are 8 edges from the vertex n to the vertex $n+1$ for each $n \geq 0$.



Now we show that

$$\log\left(R^{-1}_{E^\star_r}\right) \leq h_l(E),$$

which then implies $h_b({}^tE) = h_l(E)$ by the above proposition (hence $h_b({}^tE) \neq h_b(E)$). For a fixed vertex 0 we have

$$\begin{aligned} R^{-1}_{E^\star_r} &= \limsup_{n\to\infty} |E^n_r(0^\star)|^{1/n} \\ &= \limsup_{n\to\infty} \left|\{\alpha \in E^n_r(0) \mid s(\alpha_i) \neq 0, \text{ for } 1 \leq i \leq n\}\right|^{1/n}. \end{aligned}$$

With $n_{k+1} = 4k+1\,(k \geq 0)$,

$$|E^{n_{k+1}-1}_r(0^\star)| = |E^{4k}_r(0^\star)| = 1 + 8^{k-1} + 8^{k-4} + 8^{k-7} + \cdots,$$

and a computation gives

$$\limsup_{k\to\infty} |E^{4k}_r(0^\star)|^{\frac{1}{4k}} = 8^{1/4}.$$

But it is not hard to see that

$$\limsup_{n\to\infty} |E^n_r(0^\star)|^{\frac{1}{n}} = \limsup_{k\to\infty} |E^{4k}_r(0^\star)|^{\frac{1}{4k}},$$

hence $\log(R^{-1}_{f^\star_r}) = \log 8^{1/4} < \log 2 = h_l(E)$.

Lemma 3.4. *If E is an irreducible graph then the value*

$$\limsup_{n\to\infty} \frac{1}{n} \log |E^n(v)|$$

is independent of the choice of a vertex v.

Proof. Let v, w be two vertices of E. Then there exist two paths $\mu \in E^k$, $\nu \in E^m$ with $s(\mu) = r(\nu) = v$, $s(\nu) = r(\mu) = w$ because E is irreducible. We assume that μ and ν have the smallest length, respectively. If $\alpha = \alpha_1\alpha_2\cdots\alpha_n \in E^n(v)$ then with $i_0 = \min\{i \mid s(\alpha_i) = v\}$ write $\alpha = \alpha'\alpha''$, where $\alpha' = \alpha_1 \cdots \alpha_{i_0-1}$ and $\alpha'' = \alpha_{i_0} \cdots \alpha_n$ (if $i_0 = 1$, $\alpha = \alpha''$). Then the map

$$E^n(v) \to E^{n+k+m}(w), \quad \alpha = \alpha'\alpha'' \mapsto \alpha'\mu\nu\alpha''$$

is injective, hence $|E^n(v)| \le |E^{n+k+m}(w)|$ for each n. Therefore

$$\begin{aligned}\limsup_{n\to\infty}\frac{1}{n}\log|E^n(v)| &\le \limsup_{n\to\infty}\frac{1}{n}\log|E^{n+k+m}(w)|\\ &\le \limsup_{n\to\infty}\frac{1}{n}\log|E^n(w)|.\end{aligned}$$

□

Proposition 3.5. *Let E be an irreducible graph and $v_0 \in E^0$.*
(a) *If E is finite, then*

$$\limsup_{n\to\infty}\frac{1}{n}\log|E_l^n(v_0)| = \limsup_{n\to\infty}\frac{1}{n}\log|E^n|.$$

In particular, $h_l(E) = h_b(E) = h_b({}^tE)$.
(b) *If E is infinite, then*

$$\limsup_{n\to\infty}\frac{1}{n}\log|E^n(v_0)| \;=\; \max\{h_b(E),\, h_b({}^tE)\}.$$

Proof. (a) Let $E^0 = \{v_0, v_1, \dots, v_{k-1}\}$. Since E is irreducible there exist finite paths $\{\mu_i,\, \nu_i \mid 0 \le i \le k-1\}$ such that $s(\mu_i) = r(\nu_i) = v_0$, $r(\mu_i) = v_i = s(\nu_i)$. Suppose $|\mu_i| = m_i$, $|\nu_j| = l_j$. If $\alpha \in E^n$ is a path with $s(\alpha) = v_i$, $r(\alpha) = v_j$ then $\mu_i\alpha\nu_j \in E_l^{n+m_i+l_j}(v_0)$ is a loop at v_0. The map $\alpha \mapsto \mu_i\alpha\nu_j$ is not necessarily injective, but there exist at most k_0 paths in E^n that have the same image in $E_l^{n+m_i+l_j}(v_0)$ under the map, where $k_0 = \max_{i,j}\{m_i + l_j\}$. Hence we have

$$|E^n| \le k_0 \cdot \big|\cup_{0\le i,j\le k-1} E_l^{n+m_i+l_j}(v_0)\big| \le k_0 k^2 \max_{i,j}|E_l^{n+m_i+l_j}(v_0)|.$$

On the other hand, for each n, there exists a $k_n \in \{0, \dots, k_0\}$ such that

$$|E_l^{n+k_n}(v_0)| = \max_{i,j}|E_l^{n+m_i+l_j}(v_0)|.$$

Then $|E^n| \le k_0 k^2 |E_l^{n+k_n}(v_0)|$ and it follows that

$$\limsup_{n\to\infty}\frac{1}{n}\log|E^n| \le \limsup_{n\to\infty}\frac{1}{n}\log|E_l^n(v_0)|.$$

(b) Note first that

$$\begin{aligned}|E^n(v)| &= \big|\cup_{k=0}^n \{\alpha\beta \mid \alpha \in E_r^k(v^\star),\, \beta \in E_s^{n-k}(v_0)\}\big|\\ &= \sum_{k=0}^n \big|E_r^k(v^\star)\big|\,\big|E_s^{n-k}(v)\big| = \sum_{k=0}^n \big|({}^tE)_s^k(v^\star)\big|\,\big|E_s^{n-k}(v)\big|.\end{aligned}$$

Then

$$\begin{aligned}E(v,z) &= \sum_n \Big(\sum_{k=0}^n |({}^tE)_s^k(v^\star)|\,|E_s^{n-k}(v)|\Big) z^n\\ &= \Big(\sum_n |({}^tE)_s^n(v^\star)| z^n\Big)\Big(\sum_n |E_s^n(v)| z^n\Big)\\ &= ({}^tE)_s(v^\star, z)\cdot E_s(v, z),\end{aligned}$$

so that the radius of convergence R_E of $E(v,z)$ is equal to $\min\big\{R_{({}^tE)_s^\star},\, R_{E_s}\big\}$. Thus

$$R_E^{-1} = \max\big\{\, R_{({}^tE)_s^\star}^{-1},\, R_{E_s}^{-1}\,\big\}.$$

But Proposition 3.2 gives

$$\log\left(R^{-1}_{({}^tE)^*_s}\right) \le h_b({}^tE),$$

and also by definition $\log(R^{-1}_{E_s}) = h_b(E)$. Therefore

$$\limsup_{n\to\infty} \frac{1}{n}\log|E^n(v)| = \log(R_E^{-1}) \le \max\{h_b({}^tE),\, h_b(E)\}.$$

□

Let E be an irreducible infinite graph and let $\mathcal{D}_E$ be the commutative C^*-subalgebra of $C^*(E)$ generated by the projections $\{p_\alpha = s_\alpha s_\alpha^* \mid \alpha \in E^*\}$. Then $\mathcal{D}_E = \overline{\text{span}}\{p_\alpha \mid \alpha \in E^*\}$ and the map

$$w : \mathcal{D}_E \to C_0(X_E),\ w(p_\alpha) = \chi_{[\alpha]},$$

is a C^*-isomorphism [11]. Here $\chi_{[\alpha]}$ is the characteristic function on the cylinder set $[\alpha] = \{\beta \in X_E \mid \beta = \alpha\beta'\}$ which is both open and closed. Furthermore, from the proof of Theorem 2.4 we know that $h_{top}(\bar{X}_E) = ht(\Phi_E|_{\mathcal{D}_E})$. Put

$$\mathcal{A}_E := \overline{\text{span}}\{\, s_\alpha s_\beta^* \mid \alpha, \beta \in E^*,\ |\alpha| = |\beta|\,\}.$$

Then $\mathcal{A}_E$ is a Φ_E-invariant AF C^*-subalgebra of $C^*(E)$ with $\mathcal{D}_E \subset \mathcal{A}_E$, hence

$$ht(\Phi_E|_{\mathcal{D}_E}) \le ht(\Phi_E|_{\mathcal{A}_E}).$$

Also, for each $v \in E^0$, $\mathcal{A}_E$ contains a Φ_E-invariant AF subalgebra $\mathcal{A}_E(v)$,

$$\mathcal{A}_E(v) := \overline{\text{span}}\{\, s_\alpha s_\beta^* \mid\ r(\alpha) = r(\beta) = v,\ |\alpha| = |\beta|\,\}.$$

Lemma 3.6. *Let v be a vertex of an irreducible graph E with at least two vertices and let $n \ge 1$. Then the elements in the set*

$$\omega(n, v) = \left\{s_\alpha s_\beta^* \,\middle|\ r(\alpha) = r(\beta) = v,\ |\alpha| = |\beta| \le n\right\}$$

are linearly independent.

Proof. We prove the assertion by induction on n. For $n = 1$, suppose

$$x = \sum_{\substack{e,f\in E^1\\ r(e)=r(f)=v}} \lambda_{ef} s_e s_f^* + \lambda_0 p_v = 0.$$

If e_0 and f_0 are edges with $r(e_0) = r(f_0) = v$ and either $s(e_0) \ne v$ or $s(f_0) \ne v$ then $s_{e_0}^* p_v s_{f_0} = 0$, hence

$$0 = s_{e_0}^* x s_{f_0} = \lambda_{e_0 f_0}(s_{e_0}^* s_{e_0})(s_{f_0}^* s_{f_0}) = \lambda_{e_0 f_0} p_v,$$

thus $\lambda_{e_0 f_0} = 0$. Similarly, $\lambda_{ef} = 0$ if e and f are loops at v and $e \ne f$. Then x becomes

$$x = \sum_{e\in E_l^1(v)} \lambda_{ee} s_e s_e^* + \lambda_0 p_v = 0.$$

By irreducibility of E and the assumption that $|E^0| > 1$, there exists an edge f with $s(f) = v, r(f) \ne v$. Then $s_f s_f^* x = \lambda_0 s_f s_f^* = 0$, so that $\lambda_0 = 0$ and we have $x = \sum_{e\in E_l^1(v)} \lambda_{ee} s_e s_e^* = 0$. Since the projections $\{s_e s_e^* \mid e \in E_l^1(v)\}$ are mutually orthogonal, it follows that $\lambda_{ee} = 0$ for each $e \in E_l^1(v)$.

Now suppose that the assertion is true for $n-1$. If

$$x = \sum_{\substack{|\alpha|=|\beta|\le n \\ r(\alpha)=r(\beta)=v}} \lambda_{\alpha\beta} s_\alpha s_\beta^* = 0,\ \lambda_{\alpha\beta} \in \mathbb{C},$$

then for an edge $e \in E^1$ we have

$$0 = s_e^* x s_e = \sum_{\substack{\alpha=e\alpha' \\ \beta=e\beta'}} \lambda_{\alpha\beta} s_e^* s_\alpha s_\beta^* s_e = \sum_{|\alpha'|=|\beta'|\le n-1} \lambda_{(e\alpha')(e\beta')} s_{\alpha'} (s_{\beta'})^*.$$

Note that the elements $s_{\alpha'}(s_{\beta'})^*$ appearing in the sum are distinct. Thus by induction hypothesis, one sees that $\lambda_{(e\alpha')(e\beta')} = 0$. But the edge e was arbitrary, and so we conclude that the coefficients $\lambda_{\alpha\beta}$ are all zero. □

Proposition 3.7. (cf. [4, Proposition 2.6]) *Let $\Phi : A \to A$ be a contractive cp map of an exact C^*-algebra A. If $\{\omega_\lambda\}_{\lambda\in\Lambda}$ is a net (partially ordered by inclusion) of finite subsets in A such that the linear span of $\cup_{\lambda, l\in\mathbb{Z}^+}\Phi^l(\omega_\lambda)$ is dense in A then*

$$ht(\Phi) = \sup_\lambda ht(\Phi, \omega_\lambda).$$

Theorem 3.8. *Let E be an irreducible infinite graph. Then for each $v \in E^0$,*

$$ht(\Phi_E|_{\mathcal{A}_E(v)}) \ \le\ h_b({}^tE).$$

Proof. Let $A_n(v)$ be the C^*-subalgebra of $\mathcal{A}_E(v)$ generated by $\omega(n, v)$. Then from

$$s_\alpha s_\beta^* \cdot s_\mu s_\nu^* = \begin{cases} s_{\alpha\mu'} s_\nu^*, & \text{if } \ \mu = \beta\mu', \\ s_\alpha s_{\nu\beta'}^*, & \text{if } \beta = \mu\beta', \\ 0, & \text{otherwise,} \end{cases}$$

we see that $A_n(v) = \mathrm{span}(\omega(n, v))$ is finite dimensional.

Since $\{\omega(n, v)\}_n$ is an increasing sequence of finite subsets in $\mathcal{A}_E(v)$ such that the linear span of $\cup_n \omega(n, v)$ is dense in $\mathcal{A}_E(v)$, by Proposition 3.7 it suffices to show that

$$ht(\Phi_E, \omega(n, v)) \le h_b({}^tE), \quad n \in \mathbb{N}.$$

Set $E_l^*(v) := \cup_{k\ge 0} E_l^k(v)$ and $r(n) := |\cup_{k=0}^n E_r^k(v)|$. Fix $n_0 \in \mathbb{N}$, and define a map $\phi : \omega(n_0, v) \to M_{r(n_0)}$ by

$$\phi(s_\alpha s_\beta^*) = \sum_{\substack{|\alpha\gamma|\le n_0 \\ \gamma\in E_l^*(v)}} e_{(\alpha\gamma)(\beta\gamma)},$$

where $\{e_{\mu\nu}\}$ is the standard matrix units of of the matrix algebra $M_{r(n_0)}$. Since the elements in $\omega(n_0, v)$ are linearly independent by Lemma 3.6, one can extend the map to the linear map $\phi : A_{n_0}(v) \to M_{r(n_0)}$. Now we show that ϕ is in fact a $*$-isomorphism. To prove that it is a $*$-homomorphism, we only need to see that

$$\phi((s_\alpha s_\beta^*)(s_\mu s_\nu^*)) = \phi(s_\alpha s_\beta^*)\phi(s_\mu s_\nu^*).$$

If $\beta=\mu\beta'$, then $s_\alpha s_\beta^* s_\mu s_\nu^* = s_\alpha (s_{\nu\beta'})^*$ and

$$\begin{aligned}\phi(s_\alpha s_\beta^*)\phi(s_\mu s_\nu^*) &= \sum_{\substack{|\alpha\gamma|\le n_0\\ \gamma\in E_l^*(v)}} e_{(\alpha\gamma)(\mu\beta'\gamma)}\cdot \sum_{\substack{|\mu\delta|\le n_0\\ \delta\in E_l^*(v)}} e_{(\mu\delta)(\nu\delta)}\\ &= \sum_{\substack{|\alpha\gamma|\le n_0\\ \gamma\in E_l^*(v)}} e_{(\alpha\gamma)(\nu\beta'\gamma)} = \phi(s_\alpha(s_{\nu\beta'})^*) = \phi(s_\alpha s_\beta^* s_\mu s_\nu^*).\end{aligned}$$

If $\mu=\beta\mu'$, a similar proof works. Otherwise, $\phi((s_\alpha s_\beta^*)(s_\mu s_\nu^*))=0=\phi(s_\alpha s_\beta^*)\phi(s_\mu s_\nu^*)$. In order to show that ϕ is injective, let $\phi(\sum_{\alpha,\beta}\lambda_{\alpha\beta}\, s_\alpha s_\beta^*)=0$. Then

$$\sum_{\alpha,\beta}\lambda_{\alpha\beta}\phi(s_\alpha s_\beta^*) = \sum_{\alpha,\beta}\lambda_{\alpha\beta}\big(\sum_{\substack{|\alpha\gamma|\le n_0\\ \gamma\in E_l^*(v)}} e_{(\alpha\gamma)(\beta\gamma)}\big)=0.$$

But the vectors, $\sum_{\substack{|\alpha\gamma|\le n_0\\ \gamma\in E_l^*(v)}} e_{(\alpha\gamma)(\beta\gamma)}$ ($r(\alpha)=r(\beta)=v$, $|\alpha|=|\beta|\le n_0$), are linearly independent in $M_{r(n_0)}$. In fact, if $A:=\sum_{\alpha,\beta}\lambda_{\alpha\beta}\big(\sum_{\substack{|\alpha\gamma|\le n_0\\ \gamma\in E_l^*(v)}} e_{(\alpha\gamma)(\beta\gamma)}\big)=0$, then $e_{vv}Ae_{vv}=\lambda_{vv}e_{vv}=0$, that is, $\lambda_{vv}=0$, and for any $\alpha,\beta\in E_r^1(v)$, $e_{\alpha\alpha}Ae_{\beta\beta}=\lambda_{\alpha\beta}e_{\alpha\beta}=0$, hence $\lambda_{\alpha\beta}=0$. Repeating the process one has $\lambda_{\alpha\beta}=0$ for any $\alpha,\beta\in\cup_{k=0}^{n_0}E_r^k(v)$. Therefore $\sum_{\alpha,\beta}\lambda_{\alpha\beta}\, s_\alpha s_\beta^*=0$, and the map ϕ is injective. The surjectivity of ϕ follows from $\dim(A_{n_0}(v))=r(n_0)^2$. We simply write ϕ for $\phi: A_{n_0+l}(v)\to M_{r(n_0+l)}$ ($l\ge 0$), and $\bar\phi$ for its contractive cp extension to $\mathcal{A}_E(v)$ that exists by Arveson's extension theorem.

For each $n\in\mathbb{N}$ and $0\le l\le n-1$, note that

$$\cup_{l=0}^{n-1}\Phi_E^l(\omega(n_0,v))\subseteq \operatorname{span}(\omega(n_0+n-1,v)).$$

Then the element

$$(\bar\phi,\ \psi:=\phi^{-1},M_{r(n_0+n-1)})\in CPA(id,\mathcal{A}_E(v))$$

satisfies $\psi\circ\bar\phi|_{\omega(n_0+n-1,v)}=id_{\omega(n_0+n-1,v)}$. Thus for each $\delta>0$

$$rcp(id,\omega(n_0+n-1,v),\delta)\le r(n_0+n-1),$$

and so

$$\begin{aligned}ht(\Phi_E|_{\mathcal{A}_E(v)},\omega(n_0,v),\delta) &\le \limsup_{n\to\infty}\frac{1}{n}\log(r(n_0+n-1))\\ &=\limsup_{n\to\infty}\frac{1}{n}\log(r(n))\\ &=\limsup_{n\to\infty}\frac{1}{n}\log\big|\cup_{k=0}^{n}E_r^k(v)\big|\\ &=h_b({}^tE).\end{aligned}$$

For the last equality, note that if $k\le n$ then $\big|E_r^k(v)\big|\le\big|E_r^n(v)\big|$, hence $\big|\cup_{k=0}^n E_r^k(v)\big|\le(n+1)\cdot\big|E_r^n(v)\big|$. □

Since $h_l(E)=\sup_{E'\subset E}h(X_{E'})$ is well known (see [13, Proposition 7.2.6]), we see from Theorem 2.4 and its proof that $h_l(E)\le ht(\Phi_E|_{\mathcal{A}_E})$ holds. The following theorem gives an upper bound for $ht(\Phi_E|_{\mathcal{A}_E})$.

Theorem 3.9. *Let E be an irreducible infinite graph and $\mathcal{A}_E$ be the AF subalgebra of $C^*(E)$ generated by the partial isometries $\{s_\alpha s_\beta^* \mid \alpha,\beta\in E^*, |\alpha|=|\beta|\}$. Then*

$$ht(\Phi_E|_{\mathcal{A}_E}) \le \ \max\{h_b({}^tE), h_b(E)\}.$$

Proof. Let $E^0=\{v_1,v_2,\cdots\}$. For each $n_0\in\mathbb{N}$ and $n_1\in\mathbb{Z}^+=\{0\}\cup\mathbb{N}$, put

$$\begin{aligned}\omega(n_0,n_1) &:= \Big\{s_\alpha s_\beta^* \,\big|\, \alpha,\beta\in E^{n_1}, r(\alpha)=r(\beta)\in\{v_1,\cdots,v_{n_0}\}\Big\},\\ \omega_\Sigma(n_0,n_1) &:= \Big\{\sum s_{\alpha_i}s_{\beta_i}^* \,\big|\, s_{\alpha_i}s_{\beta_i}^*\in\omega(n_0,n_1)\Big\}.\end{aligned}$$

Note that $\omega_\Sigma(n_0,n_1)$ is not the linear span of $\omega(n_0,n_1)$. Then $\{\omega_\Sigma(n_0,n_1)\mid n_0\in\mathbb{N}, n_1\in\mathbb{Z}^+\}$ is a net of finite subsets in $\mathcal{A}_E$ which is partially ordered by inclusion. In fact, given two finite sets $\omega_\Sigma(n_0,n_1)$, $\omega_\Sigma(m_0,m_1)$ $(n_1\le m_1)$, one may write each element $s_\alpha s_\beta^*\in\omega(n_0,n_1)$ as

$$s_\alpha s_\beta^* = s_\alpha(\sum_{|\mu|=m_1-n_1} s_\mu s_\mu^*)s_\beta^* = \sum s_{\alpha\mu}(s_{\beta\mu})^*\in\omega_\Sigma(m_2,m_1),$$

where $m_2>\max\{n_0,m_0\}$ is an integer large enough so that $r(\alpha\mu)\in\{v_1,\cdots,v_{m_2}\}$ for any $\alpha\mu$ appearing in the last sum, then clearly $\omega_\Sigma(n_0,n_1)\cup\omega_\Sigma(m_0,m_1)$ is contained in $\omega_\Sigma(m_2,m_1)$.

Since the linear span of the set $\cup_{n_0,n_1,n}\Phi_E^n(\omega_\Sigma(n_0,n_1))$ is dense in $\mathcal{A}_E$, by Proposition 3.7, we show that for each finite set $\omega_\Sigma(n_0,n_1)$,

$$ht(\Phi_E,\omega_\Sigma(n_0,n_1)) \le \max\{h_b({}^tE), h_b(E)\}.$$

If $s_\alpha s_\beta^*\in\omega(n_0,n_1)$, $r(\alpha)=r(\beta)=v$, then for $l\le n-1$,

$$\Phi_E^l(s_\alpha s_\beta^*) = \sum_{|\mu|=l} s_{\mu\alpha}s_{\mu\beta}^* = \sum_{|\mu|=l} s_{\mu\alpha}(\sum_{\substack{|\nu|=n-l\\ s(\nu)=v}} s_\nu s_\nu^*)s_{\mu\beta}^* = \sum_{\substack{|\mu\alpha\nu|=n+n_1\\ |\mu|=l}} s_{\mu\alpha\nu}(s_{\mu\beta\nu})^*,$$

because $p_v=\sum_{\substack{|\nu|=n-l\\ s(\nu)=v}} s_\nu s_\nu^*$. Hence one sees that

$$\cup_{i=0}^{n-1}\Phi_E^i(\omega_\Sigma(n_0,n_1)) \subseteq \Big\{\sum_{|\mu\alpha\nu|=n+n_1} s_{\mu\alpha\nu}(s_{\mu\beta\nu})^* \,\big|\, s_\alpha s_\beta^*\in\omega(n_0,n_1)\Big\}.$$

Since the set $\{s_\mu s_\nu^*\mid \mu,\nu\in\cup_{i=1}^{n_0}E^{n_1+n}(v_i)\}$ forms a matrix units, it generates the C^*-subalgebra of $\mathcal{A}_E$ which is isomorphic to M_{k_n}, where $k_n=|\cup_{i=1}^{n_0}E^{n_1+n}(v_i)|$. Let

$$\rho_n : \mathrm{span}\{s_\alpha s_\beta^*\mid\alpha,\beta\in\cup_{i=1}^{n_0}E^{n_1+n}(v_i)\}\to M_{k_n}$$

be a $*$-isomorphism with the inverse ρ^{-1}. Then by Arveson's extension theorem ρ extends to a contractive cp map $\bar\rho:\mathcal{A}_E\to M_{k_n}$, so that we obtain an element $(\bar\rho,\rho^{-1},M_{k_n})\in CPA(id,\mathcal{A}_E)$ such that $\|\rho^{-1}\circ\bar\rho(x)-x\|=0$ if

$$x\ \in\cup_{i=0}^{n-1}\Phi_E^i(\omega_\Sigma(n_0,n_1)) \subseteq\ \mathrm{span}\{s_\alpha s_\beta^*\mid\alpha,\beta\in\cup_{i=1}^{n_0}E^{n_1+n}(v_i)\}.$$

Hence

$$rcp\big(\cup_{i=0}^{n-1}\Phi_E^i(\omega_\Sigma(n_0,n_1)),\,\delta\big)\ \le\ k_n$$

holds for any $\delta>0$. Thus

$$ht(\Phi_E,\omega_\Sigma(n_0,n_1))\ \le\ \limsup_{n\to\infty}\frac{1}{n}\log(k_n).$$

On the other hand, the irreducibility of E implies that there is an N such that $|E^{n_1+n}(v_i)| \leq |E^{n_1+n+N}(v_1)|$ for $1 \leq i \leq n_0$. Hence $k_n = |\cup_{i=1}^{n_0} E^{n_1+n}(v_i)| \leq n_0|E^{n_1+n+N}(v_1)|$. Therefore

$$\limsup_{n\to\infty} \frac{1}{n} \log k_n \leq \limsup_{n\to\infty} \frac{1}{n} \log |E^n(v_1)|,$$

and the assertion then follows from Proposition 3.5(b). □

Example 3.10. Let $E := E_{\{r_n\},\{l_n\}}$ be a Salama's infinite irreducible graph (see [18]). We assume here that $l_n + 1 \leq l_{n+1}$ for each n. There are r_k edges from the vertex $k-1$ to k, and there is only one path (of length $l_k - l_{k-1}$) from the vertex v_k to v_{k-1}.



Note that for each n, $|E^n_r(0^\star)| \leq |E^n_s(0^\star)|$, which then implies by Proposition 3.2 that

$$h_b({}^tE) \leq h_b(E).$$

Thus from Theorem 3.9, we have

$$ht(\Phi_E|_{\mathcal{A}_E}) \leq h_b(E).$$

In particular, if $E_p := E_{p,p}$ $(p > 1)$ is an irreducible infinite graph of Salama satisfying $h_l(E_p) = h_b(E_p) = \log p$, we have

$$ht(\Phi_{E_p}|_{\mathcal{A}_{E_p}}) = \log p.$$

References

[1] N. P. Brown, K. Dykema and D. Shlyakhtenko, *Topological entropy of free product automorphisms*, Acta Math. **189** (2002), 1–35.

[2] F. P. Boca and P. Goldstein, *Topological entropy for the canonical endomorphism of Cuntz-Krieger algebras*, Bull. London Math. Soc. **32** (2000), 345–352.

[3] T. Bates, D. Pask, I. Raeburn and W. Szymanski, *The C^*-algebras of row-finite graphs*, New York J. Math. **6** (2000), 307–324.

[4] N. P. Brown, *Topological entropy in exact C^*-algebras*, Math. Ann. **314** (1999), 347–367.

[5] M. Choda, *Endomorphisms of shift type (entropy for endomorphisms of Cuntz algebras)*, Operator Algebras and Quantum Field Theory (Rome, 1996), 469–475, International Press, Cambridge, MA.

[6] J. Cuntz, *Simple C^*-algebras generated by isometries*, Comm. Math. Phys. **57** (1977), 173–185.

[7] J. Cuntz and W. Krieger, *A class of C^*-algebras and topological Markov chains*, Invent. Math. **56** (1980), 251–268.

[8] V. Deaconu, *Entropy estimates for some C^*-endomorphisms*, Proc. Amer. Math. Soc. **127** (1999), no. 12, 3653–3658.

[9] K. Dykema, *Topological entropy of some automorphisms of reduced amalgamated free product C^*-algebras*, Ergodic Theory Dynam. Systems, **21** (2001), 1683–1693.

[10] B. M. Gurevic, *Topological entropy of enumerable Markov chains*, Dokl. Akad. Nauk SSSR **187** (1969), 216–226. Soviet Math. Dokl. **10** (1969), 911–915.

[11] J. A Jeong and G. H Park, *Topological entropy for the canonical conmpletely positive maps of graph C^*-algebras*, Bull. Austral. Math. Soc., to appear.

[12] E. Kirchberg, *On subalgebras of the CAR-algebra*, J. Funct. Anal. **129** (1995), no. 1, 35–63.

[13] B. P. Kitchens, *Symbolic Dynamics*, Springer 1998.

[14] A. Kumjian, Notes on C^*-algebras of graphs, Contemporary Math. 228, Operator Algebras and Operator Theory, 1998, AMS.

[15] A. Kumjian, D. Pask and I. Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. **184** (1998), 161–174.

[16] D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press 1999.

[17] C. Pinzari, Y. Watatani and K. Yonetani, *KMS states, entropy and the variational principle in full C^*-dynamical systems*, Commun. Math. Phys. **213** (2000), 331-379.

[18] I. A. Salama, *Topological entropy and recurrence of countable chains*, Pacific J. Math. **134** (1988), 325–341.

[19] E. Størmer, *Entropy of some automorphisms of the II_1 factor of the free group in infinite number of generators*, Invent. Math. **110** (1992), 63–73.

[20] D. Voiculescu, *Dynamical approximation entropies and topological entropy in operator algebras*, Comm. Math. Phys. **170** (1995), 249–281.

[21] P. Walters, *An introduction to ergodic theory*, GTM 79, Springer 1982.

MATHEMATICAL SCIENCES DIVISION, BK21, SEOUL NATIONAL UNIVERSITY, SEOUL, 151–742, KOREA, EMAIL: JAJEONG@MATH.SNU.AC.KR

DEPARTMENT OF MATHEMATICS, HANSHIN UNIVERSITY, OSAN, 447–791, KOREA, EMAIL: GHPARK-@HANSHIN.AC.KR